

## Population Genetic of Kawakawa (*Euthynnus affinis*) in Sumatera Island

Raymon Rahmanov Zedta<sup>\*1</sup>, Hawis H. Madduppa<sup>2</sup>, Neviaty Putri Zamani<sup>2</sup>, Beginer Subhan<sup>2</sup> & Lalu Mukhsin Iqbal Sani<sup>2</sup>

<sup>1</sup>Research Center for Conservation of Marine and Inland Water Resources, National Research and Innovation Agency (BRIN), Cibinong, West Java, Indonesia

<sup>2</sup>Faculty of Fisheries and Marine Science, IPB University, Bogor, West Java, Indonesia

\*Corresponding author, email: zedtaraymon@apps.ipb.ac.id

Submitted: 08 January 2023; Revised: 28 May 2024; Accepted: 03 June 2024; Published: 30 June 2024

**ABSTRACT** Over the past five decades, there has been a growing trend in the capture of kawakawa (*Euthynnus affinis*), which, alongside its potential to support small-scale commercial fisheries, highlights its significance. However, there is inadequate data on this species for suitable management and conservation status. This study aimed to investigate the genetic diversity, population structure, and connectivity of the kawakawa in the three major fishing port in Sumatera, namely Belawan, Padang, and Lampung by using DNA mitochondria control region (d-loop) Sequence. In total, 78 sequences of kawakawa with an average size of 303 bp, we found 36 polymorphic sites and 56 haplotypes from all population were identified with DNA compatibility values of 97-100%. We found the genetic diversity values in Sumatera Island were high ( $h = 0.950$ ;  $\pi = 0.034$ ), with the highest genetic diversities value in Padang ( $h = 0.990$ ;  $\pi = 0.082$ ) and the lowest in Belawan ( $h = 0.929$ ;  $\pi = 0.082$ ). AMOVA and  $F_{st}$  analyses revealed no differentiation in each population ( $F_{st} = 0.005$ ). The haplotype distribution and connectivity analyses showed genetic mixing among the three populations. This study showed a single stock at the study sites and suggests management measures at a regional level to maintain the population.

**Keywords:** Commercial fisheries; population structure; D-loop

### INTRODUCTION

Tongkol komo (*Euthynnus affinis*) (Cantor, 1849) is known in international trade as kawakawa or with the FAO code KAW (Hidayat et al., 2019). Kawakawa lives in a migratory manner in the epipelagic-neritic zone (Santos et al., 2010). This species is widely distributed in tropical and subtropical open waters (oceanic) or near coastlines (neritic) with water temperatures ranging from 18-29 °C (Collette & Nauen, 1983). Kawakawa is known for its epipelagic migratory behavior, where it migrates or swims long distances in search of food and spawning grounds (Menezes, et al., 2012). It belongs to the group of small tunas and has distinctive characteristics compared to other tuna species, such as oblique black lines curving above the lateral line and 1-4 black dots near the pectoral fin (Hidayat et al., 2018). This fish is an opportunistic predator, feeding on squid, crustaceans, mollusks, and zooplankton (Collette, 2001).

Kawakawa has high economic value in Indonesia (Amri et al., 2018). Kawakawa contributes to small-scale commercial fisheries in various countries bordering the Indian Ocean, such as India, Iran, Pakistan, and Sri Lanka. Based on data recorded by the Indian Ocean Tuna Commission (IOTC, 2020) kawakawa caught in the Indian Ocean is exploited using various fishing gear, including i.e gillnets longlines, pole-and-line, purse seine (coastal purse seine, and ring net), and other fishing gear such as baitboats, trawl, lift nets, and driftnets.

The capture trend of kawakawa in Indian Ocean has been increasing since 1950-2019. Indonesia is the top country in kawakawa catch since 2014 with average catch each year 40.000 metric ton (IOTC, 2023), followed by India, Iran, and Malaysia (IOTC, 2020). Although the results of stock assessments conducted by IOTC annually in the

past five years indicate that the kawakawa fishery is still categorized as sustainable (IOTC, 2020), there is a possibility of the stock becoming overfished and overfishing due to the increasing annual catches in the Indian Ocean. The average catch from 2014 to 2018 was 152.919 tons in the Indian Ocean. This study aims to assess the patterns of genetic population diversity and phylogenetics of several kawakawa populations in the waters of Sumatera Island. Information related to population structure in an exploited species is essential for conservation and sustainable utilization purposes. Genetic information on kawakawa in Indonesia is relatively limited, while in other countries, it is better documented, such as Malaysia, India, the Philippines, Taiwan, and Australia (Chiou & Lee 2004; Robertson et al., 2007; Santos et al. 2010; Masazurah et al., 2012; Kumar et al., 2016).

### MATERIALS AND METHODS

#### Materials

Cryogenic vials 2ml (Cat. No: 81-8204-Biologix) used for sample storage and extraction, Blue tips (Cat. No: 20-1000-Biologix) used to take up large amount of solutions up to 1000  $\mu$ L, white tips (Cat. No: 21-0010-Biologix) used to take up small amount of solutions up to 10  $\mu$ L, Yellow tips (Cat. No: 21-0200- Biologix) used to take up moderate amount of solutions up to 200  $\mu$ L, Master-Mix (My Taq TM HS Red Mix Bioline) used as pre re-prepared mixture containing all the components needed for the PCR reaction, except for the template DNA, primers, and water, D-loop Primer sequences of kawakawa (GenBank accession no.NC005318) 5'-CCGGACGTCGGAG-GTTAAAAT-3'(forward) and 5'-AGGAACCAAATGCCAGGAA-TA-3'

(reverse) (Genetika sciences) play a crucial role in initiating DNA amplification, sybergreen (Qiagen) used as a fluorescent dye that binds to double-stranded DNA (dsDNA) molecules allowing it to be used as a detection method for monitoring the amplification of the target DNA sequence in real-time, Loadingdye (Qiagen) used to help increasing density of the sample without mixing with the buffer., Agarose (Cat. No: PC0701-Vivantis) used as a matrix for separating DNA by size through electrophoresis, TAE Buffer 10X (Cat. No: V4271-Promega) used for buffer solution in agarose gel, Thermocycler gradien (Bio-rad) used to multiple annealing temperatures simultaneously across a range of temperature gradients, 96-well polypropylene deep well plate (S030754935-Corning), The samples used in this study was muscle tissue around the caudal fork excute using strile knife (Marwayana, 2015).

## Methods


### Study sites

This research was conducted from October 2019 to July 2021. Sample collection took place at several Fish Landing Sites in the waters of Sumatra Island, namely PP Belawan in North Sumatra, TPI Padang in West Sumatra, and TPI Lempasing in Lampung (Figure 1). Molecular analysis was carried out at the Oceanogen Laboratory in Bogor.

### Sample collection

A total of 78 samples of muscle tissue around the caudal peduncle (Marwayana, 2015) of kawakawa, each measuring 0.5 cm<sup>3</sup>, were taken and preserved in sample bottles containing 96% ethanol. The sample size consisted of 26 individuals from each research location. The technical procedure for muscle tissue sampling was based on a genetic sampling protocol that had been implemented through collaborative cooperation between CSIRO (Australia), AZTI Tecnalia (Spain), IRD (France), and LRPT (Indonesia).

### Extraction, Amplification, and DNA Sequencing



**Figure 1.** The map shows the locations of kawakawa sample collection at Pelabuhan Perikanan Belawan in North Sumatra, Fish Landing Site Padang in West Sumatra, and Fish Landing Site Lempasing in Lampung.

**Description:** The letter “n” in the research point indicates the number of samples taken.

The DNA fragments were extracted following the Chelex 10% DNA extraction method (Walsh et al., 2013). The DNA amplification, also known as polymerase chain reaction (PCR), targeted the half portion of the hypervariable region of the mitochondrial control region (mtDNA D-loop) using the first half of the control region (D-loop) forward primer (5' – CCG GAC GTC GG A GGT TAA AAT – 3') and the reverse primer (5' – AGG AAC CAA ATG CCA GGA ATA) (Menezes et al., 2006). Each PCR reaction in one tube had a total volume of 25 µl. The components of each reaction included 3 µl of DNA template, 12.5 µl of MasterMix (My Taq™ HS Red Mix Bionline), 7.5 µl of ddH<sub>2</sub>O, 1 µl of the forward primer, and 1 µl of the reverse primer.

The PCR profile consisted of the following steps: 90 °C for 5 minutes (denaturation), 50 °C for 1 minute (annealing), 72 °C for 1 minute (extension), and 72 °C for 5 minutes (final extension). The PCR process was repeated for 35 cycles (Menezes et al., 2006). The resulting PCR products (amplikon) were visualized using agarose gel electrophoresis with 1.5% agarose gel and ethidium bromide. Electrophoresis was carried out for 15 minutes at 200 volts and 400 mA current. The presence of DNA bands was observed using a UV transilluminator. Positive DNA results were sent for sequencing using the Sanger sequencing method (Sanger et al., 1977) with BigDye Terminator v3.1 and the Applied Biosystems 3730 protocol by 1st BASE Malaysia.

### Data analysis

A total of 77 mitochondrial DNA D-loop fragments were analyzed using the ClustalW (2.1) model (Larkin et al., 2007) and species confirmation was conducted using the Basic Local Alignment Tools (BLAST) database available in the MEGA X program (Molecular Evolutionary Genetics Analysis) (Kumar et al., 2018). The phylogenetic tree was constructed using the Neighbor-joining method with the Maximum Composite Likelihood model and 1000x bootstraps. Genetic diversity parameters, including the number of haplotypes (Hn), haplotype diversity (h), and nucleotide diversity (π), were analyzed using the DnaSP (DNA Sequence Polymorphism) 6.0 software (Rozas et al., 2017) to estimate the number of polymorphic sites and the number of unique haplotypes (nh) (Menezes, et al., 2012).


Genetic diversity analysis, haplotype diversity (h), and nucleotide diversity (Masatoshi, 1973), as well as population structure, were measured as the basis for assessing the level of genetic diversity in the mitochondrial DNA control region sequences. Population genetic structure was evaluated using the Arlequin 3.5.2.2 program (Excoffier & Lischer, 2010). The analysis of differences in genetic distances between populations was performed using the fixation index (Fst) (Excoffier et al., 1992). Genetic distances within and between populations were analyzed based on distance parameters (Nei, 1973). The distribution of haplotypes and the tree among sequences was visualized using the Network 10 application (<https://www.fluxus-engineering.com>) (Bandelt et al., 1999)

**RESULTS AND DISCUSSION**



**Figure 2.** Personal documentation of photos of kawakawa samples that were landed at TPI Belawan.

The query-based confirmation findings encompass the level of similarity (Hosein *et al.*, 2017). The 78 samples were identified as kawakawa, except for one sample that was identified as *Auxis thazard* (frigate tuna). (PDG15). This misidentification occurred due to visual errors during the sample search process in the field. Each individual kawakawa was analyzed using phylogenetic tree reconstruction, with frigate tuna/*Katsuwonus pelamis* as the outgroup (Figure 3), revealing 9 different clades. The genetic analysis of the 77 D-loop gene sequences showed a fragment DNA length of 303 base pairs (bp) from three sampling locations (Belawan, Padang, and Lampung).



**Figure 3.** The phylogenetic tree reconstruction was performed using the Maximum Composite Likelihood method with 1000x bootstraps on 77 kawakawa fish samples, with *Katsuwonus pelamis* as the outgroup.

**Table 2.** Genetic diversity of kawakawa fish, number of samples (n), number of haplotypes (Hn), genetic diversity (h), and nucleotide diversity ( $\pi$ ) at 3 research locations.

Locations	n	Genetic diversity		
		H <sub>n</sub>	h	$\pi$
Belawan	26	15	0.929±0.031	0.082±0.049
Padang	25	19	0.990±0.029	0.082±0.049
Lampung	26	22	0.963±0.027	0.080±0.048
All locations	77	46	0.950±0.016	0.034±0.020

The genetic diversity values of kawakawa populations based on haplotype (h) and nucleotide ( $\pi$ ) diversity are uniform and high (Table 2). The genetic diversity of kawakawa populations in Lampung (h = 0.963;  $\pi$  = 0.080) has the highest values, followed by Padang population (h = 0.990;  $\pi$  = 0.082), and Belawan population (h = 0.929;  $\pi$  = 0.082). Overall, the genetic diversity value of kawakawa populations in the waters of Sumatra is h = 0.950;  $\pi$  = 0.034. Research conducted in India by Kumar *et al.* (2012) on kawakawa fish showed lower genetic and nucleotide diversity values compared to those in Indonesia, which are 0.040-0.262 and 0.080-0.697, respectively.

Differences in genetic diversity values may be attributed to several factors such as the number of samples (Masatoshi, 1987), spawning patterns, random drift leading to genetic drift, distribution, and migration (Hoban *et al.*, 2021). Furthermore, Madduppa *et al.* (2021) explained that the genetic diversity value of a population is influenced by the number of haplotypes (Hn) present. A higher number of haplotypes correlates with higher genetic diversity. High haplotype diversity combined with low nucleotide diversity suggests a large population size that

has recently expanded, allowing the preservation of new alleles within the population but not enough time for additional nucleotide changes to accumulate across haplotypes (Chen *et al.*, 2014; Delrieu-Trottin *et al.*, 2017; Kasim *et al.*, 2020).

The migration of kawakawa fish also contributes to the high genetic diversity value by allowing encounters with other populations and increasing the likelihood of interbreeding between different populations. High genetic diversity values can play a role in reducing the risk of inbreeding in a population (Frankham, 2005) and providing opportunities for organisms living in their natural habitat to adapt to potential environmental changes (Wernberg *et al.*, 2018).

**Population structure**

The results of the AMOVA analysis for populations in Sumatra are presented in Table 4, and the interpopulation F<sub>st</sub> values in Table 3 indicate no significant differences among the kawakawa populations in the waters of Sumatra, with F<sub>st</sub> values ranging from 0.001 to 0.004. Fakhri *et al.* (2015) stated that F<sub>st</sub> values ≥0 and <0.5 indicate no



**Table 3.** The results of AMOVA for degrees of freedom (df), diversity (%),  $F_{st}$  values, and significance level (p) within and among kawakawa populations in Sumatra.

Variance	db	Diversity (%)	$F_{st}$	P-value
Indonesia				
Inter population	2	0.49	0.005	0.294 ± 0.013
Intra population	74	99.51		
Total	76			

significant differences, while  $F_{st}$  values  $>0.5$  and  $\leq 1$  are considered to have significant differences. These results suggest that there are no subpopulations within the kawakawa population in the waters of Sumatra.

This indication of a single stock is consistent with the findings of Kumar et al. (2012<sup>a</sup>) and Kumar et al. (2012<sup>b</sup>), who revealed a single stock of kawakawa population throughout the waters of India. The close geographic proximity allows kawakawa populations to mix during migration, increasing the likelihood of interpopulation reproductive processes in that area. These findings provide important initial evidence that kawakawa in the waters of Sumatra is still genetically connected. Similar results have been reported in studies of kawakawa population structure in Taiwan by Chiou & Lee (2004) and Filipina by Santos et al. (2010).


The high mobility of kawakawa during migration among these three populations can lead to gene flow. Kawakawa is widely distributed in open tropical and subtropical waters (oceanic) or near coastlines (neritic) with water temperatures ranging from 18-29 °C (Collette & Nauen, 1983). As a member of the genus Euthynnus, which is widely distributed in tropical and subtropical waters, kawakawa is commonly found in the Indian Ocean, Central Pacific, and Western Pacific (Ollé et al., 2021).

**Table 4.** The  $F_{st}$  values using the pairwise distance method (lower diagonal) and the significance level (p) values (upper diagonal) for kawakawa populations in Belawan, Padang, and Aceh are as follows.

Location	Belawan	Lampung	Padang
Belawan	-	0.315	0.459
Lampung	0.003	-	0.279
Padang	0.001	0.004	-

#### Population connectivity

The analysis of the distribution of 56 haplotypes from a total of 77 kawakawa samples shows connections among each kawakawa haplotype (Figure 4). The haplotypes are distributed among all individuals, making it impossible to group (clade) them based on different geographical locations. There are several connected haplotypes and others that are separate. The highest connectivity among kawakawa haplotypes is observed in haplotype 4 in Figure 4, consisting of 12 samples, indicating the mixing of individuals from all three research populations. Haplotype 4 comprises 5 individuals from the Belawan population (42.66%), 4 individuals from the Lampung population (33.33%), and 3 individuals from the Padang population (25%). Haplotype 3, on the other hand, consists of 6 samples with 4 individuals from the Belawan population and one individual each from the

**Figure 4.** Haplotype connectivity of kawakawa in the Sumatra waters.

Lampung and Padang populations.

Figure 5 illustrates the distribution of 55 haplotypes across all research locations. The Belawan population is dominated by haplotypes 3 (19%) and 4 (15%), while Padang and Lampung are dominated by haplotype 4 (12%) with only a few haplotypes 3 (4%). Haplotype 1 is found only in the Belawan population. The distribution of these haplotypes can be influenced by several factors. Firstly, it is related to the migratory nature of kawakawa species, as kawakawa is one of the pelagic fish species that migrates across oceans. The dispersal of pelagic fish larvae is influenced by ocean currents during their release into the water column (Adams et al., 2019). Haplotype and nucleotide diversity are essential indicators of genetic variation, and the variation within groups is influenced by the magnitude of these indicators (Zhang et al., 2020). Kawakawa is known for its epipelagic migratory characteristics, migrating or swimming long distances in search of food and spawning (Kumar et al., 2012).

The values of genetic diversity parameters reveal that kawakawa in Sumatra, Indonesia form a mixed population pattern, as evidenced by the wide species distribution range and low, non-significant  $F_{st}$  values across all investigated populations. Genetic analysis of kawakawa samples from three locations indicates the presence of a single stock between Belawan, Padang, and Lampung. The observed population diversity of kawakawa in the waters of Sumatra highlights the importance of developing better participatory management strategies among institutions in Indonesia, especially in Sumatra and at the regional level. A significant amount of work is needed to manage fisheries resources to preserve and restore genetic stocks for sustainable harvesting. Molecular analysis, combined with other methods, can provide a valid assessment for effective harvesting strategies (Toro et al., 2003).




Figure 5. Map of the distribution of kawakawa population haplotypes in the waters of Sumatra Island.

## CONCLUSION AND RECOMMENDATION

### Conclusion

The analysis of genetic diversity and population structure of kawakawa fish in the waters of Sumatra, Indonesia, does not show any detected population structure based on haplotype composition, indicating genetic connectivity and relatedness. Stock-based management and collaborative cross-regional management are needed to support tuna fish management efforts in Indonesia.

### Recomendation

Future research on the population genetics of kawakawa (*Euthynnus affinis*) in Sumatra should expand the sample size and geographic coverage, incorporate additional genetic markers like SNPs and mitochondrial DNA, and include temporal sampling to assess genetic changes over time. Collaborating with other research groups and integrating genetic data into conservation and management plans are essential for the sustainable management of kawakawa populations

## AUTHORS' CONTRIBUTION

RRZ is doing research idea, write manuscript and translation, data & sample collection, data analysis; HHM is doing guidance contribution, data analysis; NPZ is guidance contribution; BS is doing guidance contribution; LMIS is doing data analysis.

## ACKNOWLEDGEMENT

The author expresses gratitude to the Indonesian Endowment Fund for Education (LPDP) for providing research funding and postgraduate education scholarships. The author also thanks the Oceanogen Laboratory in Bogor, the Faculty of Fisheries and Marine Science at Bogor Ag-

ricultural University, the Department of Marine and Fisheries in Padang, West Sumatra, the Fishery Port in Belawan, North Sumatra, the fishermen at TPI Lempasing, Lampung, and the Ministry of Marine Affairs and Fisheries.

## REFERENCES

- Adams, N.L., A. Heyland, L.L. Rice & K.R. Foltz. 2019. Procuring animals and culturing of eggs and embryos. *Methods in Cell Biology*. 150: 3-46. <https://doi.org/10.1016/BS.MCB.2018.11.006>
- Amri, K., F.A. Nora, D. Emaningsih & T. Hidayat. 2018. Reproduction and spawning season of kawakawa (*Euthynnus Affinis*) based on monsoon and SST distribution in Indian Ocean South off Java-Nusa Tenggara. *Bawal Widya Riset Perikanan Tangkap*. 10 (2): 155-167.
- Bandelt, H.J., P. Forster & A. Röhl. 1999. Median-joining networks for inferring intraspecific phylogenies. *Molecular Biology and Evolution*. 16 (1): 37-48. <https://doi.org/10.1093/OXFORDJOURNALS.MOLBEV.A026036>
- Chen, W., W. Hong, S. Chen, Q. Wang & Q. Zhang. 2014. Population genetic structure and demographic history of the mudskipper *Boleophthalmus pectinirostris* on the northwestern pacific coast. *Environmental Biology of Fishes*. 98 (3): 845-856. <https://doi.org/10.1007/S10641-014-0320-1>
- Chiou, W.D & L.K. Lee. 2004. Migration of kawakawa *Euthynnus affinis* in the waters near Taiwan. *Fisheries Science*. 70 (5): 746-757. <https://doi.org/10.1111/j.1444-2906.2004.00867.x>
- Collette, B.B. 2001. Tunas (also, albacore, bonitos, mackerels, seerfishes, and wahoo). *FAO Species Identification Guide for Fishery Purposes The Living Marine Resources of the Western Central Pacific*.
- Collette, B.B & C.E. Nauen. 1983. *FAO Species Catalogue Vol . 2*

- Scombrids of the world an annotated and illustrated catalogue of Tunas, Mackerels, Bonitos and related species know to date. In FAO Fisheries Synopsis. FishMSynop. 125 (2).
- Delrieu-Trottin, E., S. Mona, J. Maynard, V. Neglia, M. Veuille & S. Planes. 2017. Population expansions dominate demographic histories of endemic and widespread Pacific reef fishes. *Scientific Reports*. 7 (1): 1-13. <https://doi.org/10.1038/srep40519>
- Excoffier, L., P.E. Smouse & J.M. Quattro. 1992. Analysis of molecular variance inferred from metric distances among DNA haplotypes: Application to human mitochondrial DNA restriction data. *Genetics*. 131 (2): 479-491. <https://doi.org/10.5962/bhl.title.86657>
- Excoffier, L & H.E.L. Lischer. 2010. Arlequin suite ver 3.5: A new series of programs to perform population genetics analyses under Linux and Windows. *Molecular Ecology Resources*. 10 (3): 564-567. <https://doi.org/10.1111/j.1755-0998.2010.02847.x>
- Fakhri, F., I. Narayani & I.G.N.K. Mahardika. 2015. Genetic diversity of skipjack tuna (*Katsuwonus pelamis*) from Jembrana and Karangasem Regencies, Bali. *Jurnal Biologi*. 19 (1). <http://garuda.ristekdikti.go.id/journal/article/366597>
- Forster, P., A. Torroni, C. Renfrew & A. Röhl. 2001. Phylogenetic star contraction applied to Asian and Papuan mtDNA evolution. *Molecular Biology and Evolution*. 18 (10): 1864-1881. <https://doi.org/10.1093/OXFORDJOURNALS.MOLBEV.A003728>
- Frankham, R. 2005. Genetics and extinction. *Biological Conservation*. 126 (2): 131-140. <https://doi.org/10.1016/J.BIOCON.2005.05.002>
- Hidayat, T., T. Nugroho & U. Chodrijah. 2019. Biologi ikan tongkol komo (*Euthynnus affinis*) di Laut Jawa. *Journal of Tropical Fisheries Management*. 2 (1): 30-36. <https://doi.org/10.29244/jppt.v2i1.25315>
- Hoban, S., C.D. Campbell, J.M. da Silva, R. Ekblom, W.C. Funk, B.A. Garner, J.A. Godoy, F. Kershaw, A.J. MacDonald, J. Mergeay, M. Minter, D. O'Brien, I.P. Vinas, S.K. Pearson, S. Pérez-Espona, K.M. Potter, I.R.M. Russo, G. Segelbacher, C. Vernesi & M.E. Hunter. 2021. Genetic diversity is considered important but interpreted narrowly in country reports to the Convention on Biological Diversity: Current actions and indicators are insufficient. *Biological Conservation*. 261: 109233. <https://doi.org/10.1016/J.BIOCON.2021.109233>
- Hosein, F.N., N. Austin, S. Maharaj, W. Johnson, L. Rostant, A.C. Ramdass & S.N. Rampersad. 2017. Utility of DNA barcoding to identify rare endemic vascular plant species in Trinidad. *Ecology and Evolution*. 7 (18): 7311-7333. <https://doi.org/10.1002/ECE3.3220>
- IOTC. 2020. Report of the 10th Session of the IOTC Working Party on Neritic Tunas | IOTC. 10th Session of the IOTC Working Party on Neritic Tunas. <https://iotc.org/documents/WPNT/10/RE>
- Kasim, N.S., T.N.A.M. Jaafar, R.M. Piah, W. Mohd Arshaad, S.A. Mohd Nor, A. Habib, M.Abd. Ghaffar, Y.Y. Sung, M. Danish-Daniel & M.P. Tan. 2020. Recent population expansion of longtail tuna *Thunnus tonggol* (Bleeker, 1851) inferred from the mitochondrial DNA markers. *PeerJ*. 8: e9679. <https://doi.org/10.7717/PEERJ.9679/SUPP-1>
- Kumar, G., M. Kocour & S.P. Kunal. 2016. Mitochondrial DNA variation and phylogenetic relationships among five tuna species based on sequencing of D-loop region. *Mitochondrial DNA. Part A, DNA Mapping, Sequencing, and Analysis*. <https://doi.org/10.3109/19401736.2014.971313>
- Kumar, G., S.P. Kunal & M.R. Menezes. 2012. Low genetic variation suggest single stock of kawakawa *Euthynnus affinis* (Cantor, 1849) along the indian coast. *Turkish Journal of Fisheries and Aquatic Sciences*. 12 (3): 555-564. [https://doi.org/10.4194/1303-2712-V12\\_3\\_02](https://doi.org/10.4194/1303-2712-V12_3_02)
- Kumar, G., S.P. Kunal, M.R. Menezes & R.M. Meena. 2012. Single genetic stock of kawakawa *Euthynnus affinis* (Cantor, 1849) along the Indian coast inferred from sequence analyses of mitochondrial DNA D-loop region. *Conservation Genetics*. 13 (4): 1119-1131. <https://doi.org/10.1007/s10592-012-0359-5>
- Kumar, S., G. Stecher, M. Li, C. Knyaz & K. Tamura. 2018. MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Molecular Biology and Evolution*. 35 (6): 1547-1549. <https://doi.org/10.1093/MOLBEV/MSY096>
- Larkin, M.A., G. Blackshields, N.P. Brown, R. Chenna, P.A. Mcgettigan, H. McWilliam, F. Valentin, I.M. Wallace, A. Wilm, R. Lopez, J.D. Thompson, T.J. Gibson & D.G. Higgins. 2007. Clustal W and clustal X version 2.0. *Bioinformatics*. 23 (21): 2947-2948. <https://doi.org/10.1093/BIOINFORMATICS/BTM404>
- Madduppa, H., R. Martaulina, Z. Zairion, R.M. Renjani, M. Kawaroe, N.P. Anggraini, B. Subhan, I. Verawati & L.M.I. Sani. 2021. Genetic population subdivision of the blue swimming crab (*Portunus pelagicus*) across Indonesia inferred from mitochondrial DNA: Implication to sustainable fishery. *PLOS ONE*, 16 (2): e0240951. <https://doi.org/10.1371/JOURNAL.PONE.0240951>
- Marwayana, O.N. 2015. Ekstraksi asam dioksiribonukleat (DNA) dari sampel jaringan otot. *Ocena* 9 (2): 1-9.
- Menezes, M.R., M. Ikeda & N. Taniguchi. 2006. Genetic variation in skipjack tuna *Katsuwonus pelamis* (L.) using PCR-RFLP analysis of the mitochondrial DNA D-loop region. *Journal of Fish Biology*. <https://doi.org/10.1111/j.0022-1112.2006.00993.x>
- Masatoshi, N. 1973. Analysis of gene diversity in subdivided populations. *Proceedings of the National Academy of Sciences of the United States of America*. <https://doi.org/10.1073/pnas.70.12.3321>
- Masatoshi, N. 1987. *Molecular Evolutionary Genetics*. Columbia University Press. <https://doi.org/10.7312/nei-92038>
- Ollé, J., L. Vilà-Valls, J. Alvarado-Bremer, G. Cerdaneres, T.Y. Duong, G. Hajjej, P.G. Lino, R. Muñoz-Lechuga, F.N. Sow, N.C. Diah, R.M. Araguas, N. Sanz & J. Viñas. 2021. Population genetics meets phylogenetics: new insights into the relationships among members of the genus *Euthynnus* (family Scombridae): *Euthynnus* population genetics and phylogeny. *Hydrobiologia*. 849 (1): 47-62. <https://doi.org/10.1007/S10750-021-04707-6/FIGURES/4>
- Polzin, T & D.S. Vahdati. 2003. On steiner trees and minimum spanning trees in hypergraphs. *Operations Research Letters*. 31 (1): 12-20. [https://doi.org/10.1016/S0167-6377\(02\)00185-2](https://doi.org/10.1016/S0167-6377(02)00185-2)
- Rozas, J., A. Ferrer-Mata, J.C. Sanchez-DelBarrio, S. Guirao-Rico, P. Librado, S.E. Ramos-Onsins & A. Sanchez-Gracia. 2017. DnaSP 6: DNA sequence polymorphism analysis of large data sets. *Molecular Biology and Evolution*. 34 (12): 3299-3302. <https://doi.org/10.1093/molbev/msx248>
- Sanger, F., S. Nicklen & A.R. Coulson. 1977. DNA sequencing with chain-terminating inhibitors. *Proceedings of the National Academy of Sciences of the United States of America*. 74 (12): 5463-5467. <https://doi.org/10.1073/pnas.74.12.5463>
- Santos, M.D., G.V. Lopez & N.C. Barut. 2010. A pilot study on the

- genetic variation of eastern little tuna (*Euthynnus affinis*) in Southeast Asia. *Philippine Journal of Science*.139 (1): 43-50.
- Tamura, K., G. Stecher & S. Kumar. 2021. MEGA11: Molecular Evolutionary Genetics Analysis version 11. *Molecular Biology and Evolution*. <https://doi.org/10.1093/molbev/msab120>
- Walsh, P.S., D.A. Metzger & R. Higushi. 2013. Chelex 100 as a medium for simple extraction of DNA for PCR-based typing from forensic material. *BioTechniques*. 54 (3): 134-139. <https://doi.org/10.2144/000114018>
- Wernberg, T., M.A. Coleman, S. Bennett, M.S. Thomsen, F. Tuya & B.P. Kelaher. 2018. Genetic diversity and kelp forest vulnerability to climatic stress. *Scientific Reports*. 8 (1). 1-8. <https://doi.org/10.1038/s41598-018-20009-9>
- Zhang, Q., C. Sun, Y. Zhu, N. Xu & H. Liu. 2020. Genetic diversity and structure of the round-tailed paradise fish (*Macropodus ocellatus*): Implications for population management. *Global Ecology and Conservation*. 21. e00876. <https://doi.org/10.1016/J.GECCO.2019.E00876>