

Accepted Manuscript

Isolation of Potential Antagonistic Rhizosphere Fungi against *Alternaria alternata* from Organic Carrot Productions

Nita Noviyanti, Susiana Purwantisari, & Agung Suprihadi

DOI : <https://doi.org/10.22146/jpti.94840>

Reference : Mak-679

To appear in : *Jurnal Perlindungan Tanaman Indonesia*

Received date : 20 March 2024

Revised date : 10 April 2024

Accepted date : 4 June 2024

This is an early version of Accepted Manuscript, which has been through the Jurnal Perlindungan Tanaman Indonesia peer-review process and is available shortly after acceptance as our service to the community. The edited manuscript will be published after technical editing, formatting, and proofreading. Please note that minor changes to the text and/or graphics might be introduced during technical editing, which could affect the content. Terms & Conditions and the Ethical Guidelines of the Journal still apply.

Research Article

Isolation of Potential Antagonistic Rhizosphere Fungi against *Alternaria alternata* from Organic Carrot Productions

Nita Noviyanti¹⁾, Susiana Purwantisari^{1)*}, & Agung Suprihadi¹⁾

¹⁾ Department of Biology, Faculty of Science and Mathematics, Diponegoro University
Jln. Prof. Soedarto SH., Semarang, Central Java 50275 Indonesia

*Corresponding author. E-mail: susiana_purwantisari@yahoo.co.id

ABSTRACT

Rhizospheric fungi can have serve as biological agents against plant pathogens *Alternaria alternata*, which infects carrot plantss. Rhizospheric fungi are more abundant in organic lands. This study aims to isolate fungi from organic carrot's rhizosphere that are able to suppress *A. alternata*'s growth and determine isolate rhizospheric fungi ability to inhibit *A. Alternata* to identify potential biological control agents against *A. alternata*. This research was conducted at the Laboratory of Biotechnology and Basic Biology, Diponegoro University, between January–June 2023. Research methods include survey location, soil sampling, rhizospheric fungi isolation, pathogenic fungi isolation, creating growth curve, pathogenicity tests, antagonism tests, and fungi identification. This study used a complete randomized design (CRD) and was analyzed using ANOVA at the 5% level of significance and Duncan's hoc-posttest. Fungi isolation from organic carrot plant's rhizosphere resulted in 13 isolates consisting of 6 genera: *Penicillium*, *Aspergillus*, *Paecilomyces*, *Myrothecium*, *Trichoderma*, and *Simplicillium*. The antagonism test showed that 8 antagonistic isolates, including the genus *Penicillium* (WO1, WO6, WO10), *Paecilomyces* (WO5), *Myrothecium* (WO7), *Trichoderma* (WO9), and *Aspergillus* (WO11, WO13), where WO1, WO6, WO10, and WO11 have a medium percentage of inhibition (44.12-57.84%), while WO5, WO7, WO9, and WO13 have a high percentage of inhibition (64.29-76.06%). The eight isolates showed antagonism mechanisms, including antibiosis (WO1, WO5, WO6, WO7, WO10, and WO11) and parasitism (WO9 and WO13).

Keywords: *Alternaria* leaf blight; antagonist fungi; antagonism test; carrot plant rhizosphere; organic soil

INTRODUCTION

Soil microorganisms, including fungi, are one of the components of soil biodiversity that play a vital role in the soil ecosystem. According to Noviyanti *et al.* (2022), fungus can act as a decomposer and play a vital role in soil fertility by providing nutrients for plants. Among other factors that influence the existence of fungi is the way land is managed. Organic land production systems that use organic materials and are based on environmentally sustainable practices. Organic land is known to have a higher abundance of soil microorganisms than conventional land (Das & Dkhar, 2011; Nurbailis *et al.*, 2014).

Carrot (*Daucus carota* L.) is one of many highly consumed vegetables because of its high nutritional content. According to Bender *et al.* (2020), organic carrots also have several advantages, such as higher market-demand for organic products, low nitrate content, high vitamin C and beta-carotene content, as well as lower chemical residue levels. Based on Soesanto *et al.* (2013), pathogens are challenges in plant production because they can reduce yield production. An example of carrot pathogen is *Alternaria alternata*, which causes leaf and plant blight (Soyal *et al.*, 2018; Gayithri *et al.*, 2021; Yadav *et al.*, 2021). Control of *A. alternata* has been largely done with the use of synthetic fungicides. The use of synthetic fungicides cause harm to the environment and humans. Thus, environmental friendly disease management are needed, for example, using antagonistic fungi (Muksin *et al.*, 2013). Rhizospheric fungi that are abundant in organic soils are opportunities to identify potential biocontrol agents against pathogens that attack carrot plants, including *A. alternata*. The objectives of this study include examining the antagonism ability of fungi isolated from the rhizosphere of organic carrot plants against *A. alternata*, examining the macroscopic and microscopic characteristics of fungi from the rhizosphere of organic carrot, and determine types of fungi with potential to suppress the growth of *A. alternata*.

MATERIALS AND METHODS

Research Location. The soil sampling from organic land was done at “Argo Ayuning Tani” farmer group in Senden Village, Selo District, Boyolali Regency (7.4849918, 110.4716418). The research was carried out at Biotechnology and Basic Biology Laboratory, Biology Department, Faculty of Science and Mathematics, Diponegoro University.

Soil Sampling. Soil was collected from areas directly in contact with healthy carrot roots from 3 plant samples, were randomly selected, and combined into one as a composite sample (Purwantisari *et al.*, 2016). The sample was stored in a cool box (18 x 28 x 10 cm) and brought to the laboratory for further isolation (Amaria *et al.*, 2013).

Isolation and Purification of Rhizospheric Fungi. A total of 10 g of soil sample was dissolved in 90 mL of sterile distilled water and homogenized using a shaker at 120 rpm for 15 minutes at room temperature (25–30 °C). One milliliter was transferred to a test, which contained 9 mL of distilled water, homogenized to obtain 10⁻¹ dilution. Dilutions were repeated until 10⁻⁵ dilution was obtained. One milliliter of the suspension resulting from 10⁻¹–10⁻⁵ dilution was taken and transferred aseptically to a petri dish containing Potato Dextrose Agar (PDA) medium (Himedia, India) added with chloramphenicol. Fungi was incubated for

3–5 days at room temperature (Yulia *et al.*, 2022). Fungi grown from a 10^{-5} dilution were purified based on the appearance of the colonies by transferring each colony to a new sterile PDA.

Preparation of Growth Curves. Growth of each purified isolate grown using the single dot method was observed. Colony diameter was measured every day until the 15th day. Then a growth curve is made from these results (Hasanah, 2018).

Isolation of Pathogenic Fungi. Symptomatic carrot plants were sampled and infected tissues were directly plated. Samples were washed thoroughly using water, then cut into 1×1 cm pieces using a scalpel. The sample surface was then sterilized by immersion in a 70% alcohol solution for 30 seconds and NaOCl 1% for 1 minute, then rinsed using sterile distilled water. Furthermore, the sample pieces were dried on sterile tissue paper and planted on sterile PDA medium that had been treated with chloramphenicol. Then, the culture was incubated for 2–5 days at room temperature (Asrul *et al.*, 2021). The growing fungi were purified for use in the pathogenicity test, and identification of the pathogenic fungi was carried out based on their macroscopic and microscopic characteristics.

Pathogenicity Test. Healthy carrot tubers were washed thoroughly using water and cut into ± 5 mm thick pieces (Tülek & Dolar, 2015). The tuber pieces were then sterilized by immersion in 70% alcohol for 1 minute and 1% NaOCl for 1 minute, then rinsed using sterile distilled water three times (Fadhilah *et al.*, 2014). The tuber pieces were placed on a sterile petri dish that has been given a piece of sterile filter paper that is moistened with sterile distilled water. The tuber sections were then injured and inoculated with 5 mm mycelial-plugs of rejuvenated pathogenic cultures. Control was inoculated using agar on tuber pieces (Tülek & Dolar, 2015). The cultures were then incubated for 7 days, and symptoms were observed.

Antagonism Test. Rhizospheric and pathogenic fungi at the same growth phase (log phase) were inoculated on Petridishes (9 cm of diameter) containing PDA medium at a distance of 3 cm. Incubated for 7 days at room temperature and radius of the pathogen in the treatment and control were measured to then calculate the percentage of inhibition (%) using the the following equation (1) (Živković *et al.*, 2010):

$$P = \frac{(r1 - r2)}{r1} \times 100\% \quad (1)$$

P: inhibition percentage (%)

r1: pathogen colony radius in control

r₂: pathogen colony radius in treatment

Observation of antibiosis can be seen from the presences of a clear zone between two fungi coloy (Nurbailis *et al.*, 2014). While antagonism test that showed high inhibition, contact with hyphae, and the invasion by rhizosphere fungi, were further observed for their parasitism ability using the slide culture method to observed penetration, coiling, or lysis of pathogenic hypha under a microscope (Nurbailis *et al.*, 2014).

Identification. This was done by observing the macroscopic and microscopic characteristics of the isolates. Observed characteristics were matched with several identification books, i.e. *Pictorial Atlas of Soil and Seed Fungi Morphologies of Cultured Fungi and Key to Species* (2nd Edition) by Watanabe (2002), *Illustrated Genera of Imperfect Fungi* (4th Edition) by Barnett and Hunter (1998), *Illustrated Genera of Imperfect Fungi* (2nd Edition) by Barnett (1960), and *Food and Indoor Fungi* by Samson *et al.* (2010).

Data Analysis. The antagonism test was carried out using a completely randomized design (CRD) with three replications for each treatment. The number of treatments carried out was same as isolated rhizosphere fungi. Data were then analyzed for variance using a 5% ANOVA and Duncan's further test.

RESULTS AND DISCUSSION

Rhizospheric fungi isolation from organic carrot plants resulted in 13 types of fungi (Table 1). Pathogenic fungi isolated from symptomatic carrot plants was identified as *A. alternata* (Figure 1). This fungus showed leaf blight on isolated carrot plants. According to Soyol *et al.* (2018), Chrapačienė *et al.* (2021), Gayithri *et al.* (2021), Shebl *et al.* (2021), and Yadav *et al.* (2021), all confirmed that *A. alternata* causes leaf blight in carrots.

Based on the results of the *in vitro* pathogenicity test, pathogenic fugal isolates caused black rot on the tuber pieces. Based on previous research by Shebl *et al.* (2021) and Tülek & Dolar (2015), an *in vitro* pathogenicity test can be conducted on carrots. Although *A. alternata* is a leaf blight pathogen, it still showed root rot symptoms. This is in accordance with Shebl *et al.* (2021), who also conducted a pathogenicity test on carrot tubers and resulted in tuber rot caused the growing leaves to have lesions and then wither. This was also stated by Chrapačienė *et al.* (2021) and Yadav *et al.* (2021), *A. alternata* can also cause root rot in carrots.

Fungal growth consists of several stages; this is in accordance with Prayitno and Hidayati (2017), who stated that fungal growth consists of an adaptation phase, an

exponential phase, a stationary phase, and a death/autolysis/cell damage phase. The results of measuring the diameter of each isolate of rhizosphere fungi and pathogenic fungi show different growth rates (Figure 2). Speed of fungal growth results based on daily diameter increase, successively from the fastest resulted in: WO9, WO13, PAA, WO4, WO8, WO3, WO7, WO1, WO5, WO12, WO2, WO11, WO6, and WO10. Highest growth rates for was shown by WO9 (*Trichoderma* sp.) and WO13 (*Aspergillus* sp. 5) demonstrating growth rates faster than *A. alternata* by 4-folds while other isolates grew slower than *A. alternata*. Fungi with high growth rates and antagonistic mechanisms such as antibiotics and parasitism are potential biocontrol agents. According to Halwiyah *et al.* (2019), fungi with high growth rates are able to absorb nutrients and compete with pathogens, but slow growing fungi can still be potential biocontrol agents when they show antagonistic mechanisms such as antibiotics and/or parasitism.

The antagonistic test of rhizosphere fungi against *A. alternata* consisted of 13 treatments (Figure 3). Analysis of variance (ANOVA) showed that the inhibition percentage of antagonistic fungi significantly affected *A. alternata* in vitro at the 7th day (Table 2). Inhibition percentage of isolate WO1 (*Penicillium* sp. 1) was not significantly different from WO2 (*Penicillium* sp. 2) and WO3 (*Aspergillus* sp. 1), where all three had an inhibition percentage of 48.57%. The inhibition percentage of isolate WO4 (*Aspergillus* sp. 2) was 57.14% and not significantly different from isolate WO8 (*Aspergillus* sp. 3) that was 57.14%, isolate WO10 (*Penicillium* sp. 4) was 57.84%, and isolate WO12 (*Simplicillium* sp.) was 55.88%. The inhibition percentage of isolate WO5 (*Paecilomyces* sp.) was 64.29% and not significantly different from isolate WO7 (*Myrothecium* sp.) which was 65.20%. The inhibition percentage of isolate WO9 (*Trichoderma* sp.) was not significantly different from isolate WO13 (*Aspergillus* sp. 5), both of which were 76.06%. While the inhibition percentage of the WO6 isolate (*Penicillium* sp. 3) was 44.12%, and WO11 isolate (*Aspergillus* sp. 4) was 50.00%, both were significantly different from other rhizosphere fungi isolates.

According to Yulia *et al.* (2022), inhibition ability by antagonistic fungi are categorized into low (0–29%), moderate (30–59%), and high (60–100%). Isolates with a high inhibition percentage, included WO5 (*Paecilomyces* sp.) was 64.29%, WO7 (*Myrothecium* sp.) was 65.20%, WO9 (*Trichoderma* sp.), and WO13 (*Aspergillus* sp.5) were 76.06%. Meanwhile other isolates have a moderate percentage of inhibition, including WO1 (*Penicillium* sp. 1), WO2 (*Penicillium* sp. 2), WO3 (*Aspergillus* sp. 1) were 48.57%; WO4 (*Aspergillus* sp. 2) and WO8 (*Aspergillus* sp. 3) were 57.14%; WO6 (*Penicillium* sp. 3) was 44.12%; WO10

(*Penicillium* sp. 4) was 57.84%; WO11 (*Aspergillus* sp. 4) was 50.00%; and WO12 (*Simplicillium* sp.) was 55.88%.

Antagonism tests against *A. alternata* pathogenic fungi showed several different mechanisms (Table 3). All isolates showed a competition mechanism, but among the 13 isolates, there were isolates that showed an antibiosis mechanism, including WO1 (*Penicillium* sp. 1), WO5 (*Paecilomyces* sp.), WO6 (*Penicillium* sp. 3), WO7 (*Myrothecium* sp.), WO10 (*Penicillium* sp. 4), and WO11 (*Aspergillus* sp. 4). This was indicated by the clear zone between the antagonistic isolate and the pathogen isolate indicating that the six isolates produce secondary metabolites that inhibit the growth of the pathogen. Based on Khokhar *et al.* (2011) and Gupta and Rodriguez-Couto (2017) *Penicillium* sp. can produce compounds that act as antifungals, namely formamidine, uvidin, thymol, camazulene, altenusin, dehydroaltenusin, and fatty acids. Moreno-Gavira *et al.* (2020) and Moreno-Gavira *et al.* (2021) stated that *Paecilomyces* sp. can produce secondary metabolites as antimicrobials, including antifungals, 6-decyl salicylic acid, eicosenoic acid, ascofuranone, botulin, paecilaminol, paeciloxazine, etc. Based on Elkhateeb and Daba (2019) and Xiong *et al.* (2021), *Myrothecium* sp. can produce compounds in the form of myrothins AF, which act as antifungals against several pathogenic fungi that cause root rot; verrucarins and roridins, which act as mycotoxins; and triterpene glycosides, which inhibit pathogenic fungi in vitro. Whereas WO11 (*Aspergillus* sp. 4), which belongs to the subgenus *Polypaecilum*, mentioned by Pitt and Hocking (2022) that *Polypaecilum* does not produce harmful mycotoxins. Also, research conducted by Sabariah (2002) found that *Polypaecilum* can produce metabolites that inhibit aflatoxin production by *A. parasiticus* and *A. flavus*; these metabolites include -1,4 glycosides, L-amino acids, and lysyl or argynyl residues.

While WO9 (*Trichoderma* sp.) and WO13 (*Aspergillus* sp. 5) were capable of invading or growing against pathogenic fungi in dual culture (Figure 3), WO9 and WO13 isolates have the highest inhibition activity and were capable of invading *A. alternata*. So, we observed the mechanism of parasitism with the slide culture method and showed the results under the microscope in Figure 4.

The mechanism of parasitism by WO9 (*Trichoderma* sp.) isolates in the form of implantation, followed by coiling (a1) where the antagonist hypha envelopes the pathogen hypha, then the penetration (a2) of antagonistic hypha penetrates into the lumen of the hypha cell pathogen, as well as the enzyme degradation of the cell hypha fungus pathogen by antagonistic fungus (a3). The same is true of WO13 (*Aspergillus* sp.5) which shows an implantation that is followed by enlargement (b1), then penetration (b2) by antagonistic

fungus hypha into pathogenic cell hypha lumen with the help of cell wall degradation enzymes, as well as the enlightenment (b3) of the pathogen hypha by the antagonist fungus. This is in line with Amaresan *et al.* (2020), which stated that mycoparasitism is a complex mechanism consisting of several stages, including identification and chemotropism, implantation and depilation, secretion of extra-cellular enzymes, penetration of cell walls by hypha, and degradation of the host hypha cell wall.

The process of penetration and smoothing is due to the presence of cell wall degradation enzymes. *Trichoderma* and *Aspergillus* are known to have cell wall-degradation enzymes that play a role in parasitism. According to Herrera-Estrella (2014), *Trichoderma* sp. can produce the enzymes chitinase, glucanase, and protease involved in mycoparasitism. As Hu *et al.* (2014) mentioned, *Aspergillus* sp. could be mycoparasite with the production of chitinase and glucanase. WO9 and WO13 isolates can both produce cell wall degradation enzymes, especially chitinase, but they resist or do not experience self-degradation because they have genes that can encode cell wall protective proteins. It is consistent with Hu *et al.* (2014) and Gruber & Seidl-Seiboth (2012), which mention that chitinase can be produced by some species of *Aspergillus* spp., which has a protein-coding gene that protects the hydrophobic cell wall, similar to that found in *Trichoderma* spp.

CONCLUSION

Rhizospheric fungi isolated from organic carrot plants resulted in 13 species that have different characteristics. The fungi found belonged to six different genera: *Penicillium* sp., *Aspergillus* sp., *Paecilomyces* sp., *Myrothecium* sp., *Trichoderma* sp., and *Simplicillium* sp. The antagonistic abilities were showed by isolate WO1 (*Penicillium* sp. 1), WO5 (*Paecilomyces* sp.), WO6 (*Penicillium* sp. 3), WO7 (*Myrothecium* sp.), WO9 (*Trichoderma* sp.), WO10 (*Penicillium* sp. 4), WO11 (*Aspergillus* sp.4), and WO13 (*Aspergillus* sp. 5). WO1, WO6, WO10, and WO11 have moderate inhibition (44.12-57.84%) with antibiotic mechanism, whereas WO5, WO7, WO9, and WO13 isolates have high inhibitory percentages (64.29-76.06%) and show antibiosis mechanisms by WO5 and WO7, as well as parasitism by WO9 and WO13. The eight isolates could potentially be used as biocontrol agents if isolate a later proven or certified as non-pathogenic to host plants before application to plants.

ACKNOWLEDGEMENT/S

We thank the Department of Biology, Faculty of Sciences and Mathematics, Diponegoro University for the facilities.

LITERATURE CITED

Amaresan, N., Kumar, M.S., Annapurna, K., Kumar, K., & Sankaranarayanan, A. (2020). *Beneficial Microbes in Agro-Ecology*. Amsterdam: Elsevier.

Amaria, W., Taufiq, E., & Harni, R. (2013). Seleksi dan Identifikasi Jamur Antagonis sebagai Agens Hayati Jamur Akar Putih *Rigidiporus Microporus* pada Tanaman Karet [Selection and Identification of Antagonistic Fungi as Biological Agents of White Root Disease (*Rigidiporus microporus*) in Rubber]. *Buletin RISTRI*, 4(1), 55–64. Retrieved from <https://repository.pertanian.go.id/server/api/core/bitstreams/e8714b9a-33b9-4964-8ddd-4e557e051752/content>

Asrul, Rosmini, Rista, A., Astuti, I.D., & Yulianto, A. (2021). Karakterisasi Jamur Penyebab Penyakit Busuk Pangkal Batang (*Basal Rot*) pada Bawang Wakegi (*Allium x wakegi* Araki) [Characterization of Fungus Causing Basal Rot Disease on Wakegi Onions (*Allium x wakegi* Araki)]. *Agro Bali: Agricultural Journal*, 4(3), 341–350. <https://doi.org/10.37637/ab.v4i3.835>

Barnett, H.L. (1960). *Illustrated Genera of Imperfect Fungi*. 2nd Edition. Minnesota: Burgess Publishing Company.

Barnett, H.L. & Hunter, B.B. (1998). *Illustrated Genera of Imperfect Fungi*. 4th Edition. Minnesota: The American Phytopathological Society.

Bender, I., Edesi, L., Hiiesalu, I., Ingver, A., Kaart, T., Kaldmäe, H., Talve, T., Tamm, I., & Luik, A. (2020). Organic Carrot (*Daucus carota* L.) Production Has an Advantage over Conventional in Quantity as Well as in Quality. *Agronomy*, 10(9), 1420. <https://doi.org/10.3390/agronomy10091420>

Chrapaćienė, S., Rasiukevičiūtė, N., & Valiuškaitė, A. (2021). Biocontrol of Carrot Disease-Causing Pathogens Using Essential Oils. *Plants*, 10(11), 2231. <https://doi.org/10.3390/plants10112231>

Das, B.B., & Dkhar, M.S. (2011). Rhizosphere Microbial Population and Physico-Chemical Properties as Affected by Organic and Inorganic Farming Practices. *American-Eurasian Journal of Agricultural & Environmental Science*, 10(2), 140–150. Retrieved from [https://idosi.org/aejaes/jaes10\(2\)/2.pdf](https://idosi.org/aejaes/jaes10(2)/2.pdf)

Elkhateeb, W.A., & Daba G.M. (2019). *Myrothecium* as Promising Model For Biotechnological Applications, Potentials and Challenges. *Biomedical Journal of Scientific & Technical Research*, 16(3), 12126–12131. <https://doi.org/10.26717/BJSTR.2019.16.002869>

Fadhilah, S., Wiyono, S., & Surahman, M. (2014). Pengembangan Teknik Deteksi Fusarium Pathogen pada Umbi Benih Bawang Merah (*Allium ascalonicum*) di Laboratorium [Development of Detection Technique for Fusarium Pathogen on Seedling Shallot (*Allium ascalonicum*) Bulb at Laboratory]. *Jurnal Hortikultura*, 24(2), 171–178. Retrieved from <https://media.neliti.com/media/publications/82459-none-8b9a5458.pdf>

Gayithri, M., Ahir, R.R., Yadav, P.D., & Meena, R. (2021). Effect of Physical Parameters on The Growth of *Alternaria Alternata* Causing Alternaria Leaf Blight of Carrot. *Pharma Innovation*, 10(11), 1078–1083. Retrieved from <https://www.thepharmajournal.com/archives/?year=2021&vol=10&issue=11&ArticleId=8912>

Gruber, S., & Seidl-Seiboth, V. (2012). Self Versus Non-Self: Fungal Cell Wall Degradation in *Trichoderma*. *Microbiology*, 158(1), 26–34. <https://doi.org/10.1099/mic.0.052613-0>

Gupta, V.K., & Rodriguez-Couto, S. (2017). *New and Future Developments in Microbial Biotechnology and Bioengineering*. Amsterdam: Elsevier.

Halwiyah, N., Ferniah, R.S., Raharjo, B., & Purwantisari, S. (2019). Uji Antagonisme Jamur Patogen *Fusarium solani* Penyebab Penyakit Layu pada Tanaman Cabai dengan Menggunakan *Beauveria bassiana* Secara *In Vitro*. *Jurnal Akademika Biologi*, 8(2), 8–17. Retrieved from <https://ejournal3.undip.ac.id/index.php/biologi/article/view/24754>

Hasanah, U. (2018). Kurva Pertumbuhan Jamur Endofit Antijamur Candida dari Tumbuhan Raru (*Cotylelobium melanoxylon*) Genus *Aspergillus* [Growth Curve Endophytic Antifungal Fungi Candida of Plants Raru (*Cotylelobium melanoxylon*) *Aspergillus* Genus]. *Jurnal Biosains (Journal of Biosciences)*, 4(2), 102–107. <https://doi.org/10.24114/jbio.v4i2.10474>

Herrera-Estrella, A. (2014). Chapter 33-Genome Wide Approaches toward Understanding Mycotrophic *Trichoderma* Species. In V.K. Gupta, M. Schmoll, A. Herrera-Estrella, R.S. Upadhyay, I. Druzhinina, & M.G. Tuohy (Eds.), *Biotechnology and Biology of Trichoderma*, (pp. 455–464). Elsevier. <https://doi.org/10.1016/B978-0-444-59576-8.00033-3>

Hu, X., Webster, G., Xie, L., Yu, C., Li, Y., & Liao, X. (2014). A New Mycoparasite, *Aspergillus* sp. ASP-4, Parasitizes the Sclerotia of *Sclerotinia sclerotiorum*. *Crop Protection*, 54, 15–22. <https://doi.org/10.1016/j.cropro.2013.07.014>

Khokhar, I., Mukhtar, I., & Mushtaq, S. (2011). Antifungal Effect of *Penicillium* Metabolites against Some Fungi. *Archives of Phytopathology and Plant Protection*, 44(14), 1347–1351. <https://doi.org/10.1080/03235408.2010.496584>

Moreno-Gavira, A., Diáñez, F., Sánchez-Montesinos, B., & Santos, M. (2021). Biocontrol Effects of *Paecilomyces variotii* against Fungal Plant Diseases. *Journal of Fungi (Basel)*, 7(6), 415. <https://doi.org/10.3390/jof7060415>

Moreno-Gavira, A., Huertas, V., Diáñez, F., Sánchez-Montesinos, B., & Santos, M. (2020). *Paecilomyces* and Its Importance in the Biological Control of Agricultural Pests and Diseases. *Plants (Basel)*, 9(12), 1746. <https://doi.org/10.3390/plants9121746>

Muksin, R., Rosmini., & Panggeso, J. (2013). Uji Antagonisme *Trichoderma* sp. terhadap Jamur Patogen *Alternaria porri* Penyebab Penyakit Bercak Daun Ungu pada Bawang Merah Secara *In Vitro* [Antagonism Test *Trichoderma* sp. against Fungi Pathogens Causing Spot Disease *Alternaria porri* Purple on Shallot In-Vitro]. *e-J. Agrotekbis*, 1(2), 140–144. Retrieved from <https://media.neliti.com/media/publications/246702-uji-antagonisme-trichoderma-sp-terhadap-bdb4e753.pdf>

Noviyanti, N., Mahardhika, W.A., Lunggani, A.T., & Putra, I.P. (2022). Macrofungi Inventory at the Pine Forest of Kragilan, Magelang. *Biovalentia: Biological Research Journal*, 8(2), 138–144. <https://doi.org/10.24233/biov.8.2.2022.248>

Nurbailis, Martinius, & Azniza, V. (2014). Keanekaragaman Jamur pada Rhizosfer Tanaman Cabai System Konvensional dan Organic dan Potensinya sebagai Agen Pengendali Hayati *Colletotrichum gloeosporoides* [Fungal Diversity of Chili Rhizosphere at Conventional and Organic Cropping System and Its Role as Biocontrol Agent of *Colletotrichum gloeosporoides*]. *Jurnal Hama dan Penyakit Tumbuhan Tropika (Journal of Tropical Plant Pests and Diseases)*, 14(1), 16–24. Retrieved from <https://jhpttropika.fp.unila.ac.id/index.php/jhpttropika/article/view/306>

Pitt, J.I., & Hocking, A.D. (2022). *Fungi and Food Spoilage*. 4th Edition. Cham, Switzerland: Springer Nature.

Prayitno, T.A., & Hidayati, N. (2017). *Pengantar Mikrobiologi*. Malang, Indonesia: Media Nusa Creative.

Purwantisari, S., Priyatmojo, A., Sancayaningsih, R.P., & Kasiamdari, R.S. (2016). Penapisan Cendawan *Trichoderma* spp. untuk Pengendalian *Phytophthora infestans* secara *In Vitro* [*In Vitro* Screening of *Trichoderma* spp. as Biocontrol Agents of *Phytophthora infestans*]. *Jurnal Fitopatologi Indonesia*, 12(3), 96–103. Retrieved from <https://journal.ipb.ac.id/index.php/jfiti/article/view/12740>

Sabariah, V. (2002). *Potential Biological Control of Aflatoxins in Dried Fish*. [Thesis]. Tasmania, Australia: University of Tasmania.

Samson, R.A., Houbraken, J., Thrane, U., Frisvad, J.C., & Andersen, B. (2010). *Food and Indoor Fungi*. (CBS Laboratory Manual Series). Utrecht, The Netherlands: CBS-KNAW Fungal Biodiversity Centre.

Shebl, A.M., El-Gobashy, S.F., Arafa, R.A., Kamel, S.M., & Essa, T.A. (2021). First Record of Carrot Leaf Blight Caused by *Alternaria Alternata* in Egypt. *Egyptian Journal of Phytopathology*, 49(2), 150–151. <https://doi.org/10.21608/ejp.2021.105202.1046>

Soesanto, L., Mugiastuti, E., Rahayuniati, R.F., & Dewi, R.S. (2013). Uji Kesesuaian Empat Isolate *Trichoderma* spp. dan Daya Hambat *In Vitro* terhadap Beberapa Patogen Tanaman [Compatibility Test of Four *Trichoderma* spp. Isolates and *In Vitro* Inhibition Ability on Several Plant Pathogens]. *Jurnal Hama dan Penyakit Tumbuhan Tropika (Journal of Tropical Plant Pests and Diseases)*, 13(2), 117–123. <https://doi.org/10.23960/j.hptt.213117-123>

Soyal, S., Ghasolia, R.P., & Sarita. (2018). Histopathological Studies of *Alternaria alternata* Associated with Carrot Seed, *Daucus carota* L. *Journal of Pharmacognosy and Phytochemistry*, 7(4), 866–868. Retrieved from <https://www.phytojournal.com/archives/2018.v7.i4.5030/histopathological-studies-of-Itemgtalternaria-alternata-ltemgtassociated-with-carrot-seed-ltemgtdaucus-carota-ltemgtl>

Tülek, S., & Dolar, F.S. (2015). Detection and Identification of *Alternaria* Species Causing Diseases of Carrot in Ankara Province, Turkey. *Horticulture*, 59, 263–268. Retrieved from <https://horticulturejournal.usamv.ro/pdf/2015/art42.pdf>

Watanabe, T. (2002). *Pictorial Atlas of Soil and Seed Fungi: Morphologies of Cultured Fungi and Key to Species*. 2nd Edition. Boca Raton, Florida: CRC Press. <https://doi.org/10.1201/9781420040821>

Xiong, D.S., Yang, Y.B., Hu, B.Y., Miao, C.P., Wang, Y.L., Zou, J.M., Li, L., Li, Y.Q., Luo, X.D., & Zhao, L.X. (2021). Myrothins A–F from Endophytic Fungus *Myrothecium* sp. BS-31 Harbored in *Panax notoginseng*. *Chemistry & Biodiversity*, 18(3), e2000964. <https://doi.org/10.1002/cbdv.202000964>

Yadav, P., Choudhary, A., Verma, J., Ram, D., & Yadav, P.D. (2021). Assessment of Disease Incidence of Leaf Blight (*Alternaria alternata*) Disease of Carrot in Jodhpur District of Rajasthan. *Agrica*, 10(2), 135–139. <https://doi.org/10.5958/2394-448X.2021.00020.1>

Yulia, E., Rahayu, A., & Suganda, T. (2022). Antagonisme Jamur Rhizosfer Tanaman Karet terhadap *Rigidoporus microsporus* Secara *In Vitro* dan *In Planta* [Antagonism of Rubber Plant Rhizospheric Fungi against *Rigidoporus microsporus* in *In Vitro* and *In Planta*]. *Jurnal Agro*, 9(1), 64–79. <https://doi.org/10.15575/17824>

Živković, S., Stojanović, S., Ivanović, Z., Gavrilović, V., Popović, T., & Balaž, J. (2010). Screening of Antagonistic Activity of Microorganisms against *Colletotrichum acutatum* and *Colletotrichum gloeosporioides*. *Archives of Biological Sciences*, 62(3), 611–623. <https://doi.org/10.2298/ABS1003611Z>

APPENDIX

Table

Table 1. Identification of rhizosphere fungi isolated from organic plants

No	Isolate code	Fungi genus
1	WO1	<i>Penicillium</i> sp. 1
2	WO2	<i>Penicillium</i> sp. 2
3	WO3	<i>Aspergillus</i> sp. 1
4	WO4	<i>Aspergillus</i> sp. 2
5	WO5	<i>Paecilomyces</i> sp.
6	WO6	<i>Penicillium</i> sp. 3
7	WO7	<i>Myrothecium</i> sp.
8	WO8	<i>Aspergillus</i> sp. 3
9	WO9	<i>Trichoderma</i> sp.
10	WO10	<i>Penicillium</i> sp. 4
11	WO11	<i>Aspergillus</i> sp. 4
12	WO12	<i>Simplicillium</i> sp.
13	WO13	<i>Aspergillus</i> sp. 5

Table 2. Antagonism test of rhizosphere fungi against *Alternaria alternata* on the 7th day

Treatment/isolate	Inhibition Percentage (%)
WO1	48.57 ab
WO2	48.57 ab
WO3	48.57 ab
WO4	57.14 c
WO5	64.29 d
WO6	44.12 a
WO7	65.20 d
WO8	57.14 c
WO9	76.06 e
WO10	57.84 c
WO11	50.00 b
WO12	55.88 c
WO13	76.06 e

Note: Means followed by the same lowercase alphabet in the same column were not significantly different based on Duncan's multiple range test at the level of 5 %.

Table 3. Antagonism mechanisms of rhizosphere fungi isolates on *Alternaria alternata*

Treatment	Mechanism		
	Competition	Antibiosis	Parasitism
WO1 X PAA	+	+	-
WO2 X PAA	+	-	-
WO3 X PAA	+	-	-
WO4 X PAA	+	-	-
WO5 X PAA	+	+	-
WO6 X PAA	+	+	-
WO7 X PAA	+	+	-
WO8 X PAA	+	-	-
WO9 X PAA	+	-	+
WO10 X PAA	+	+	-
WO11 X PAA	+	+	-
WO12 X PAA	+	-	-
WO13 X PAA	+	-	+

Note: (+) indicates a mechanism; (-) indicates no mechanism; PAA (Pathogen *Alternaria alternata*)

Charts




Figure 2. Growth curve of carrot rhizosphere fungi from organic fields (WO1 to WO13) and pathogenic fungi (PAA) in PDA.

Images




Figure 1. Characteristics of *Alternaria alternata*. a. Upper colony, b. Reverse colony, c, d, e, f, g. are microscopic features(1: hyphae insulated; 2: conidiophores; 3: series of; 4: conidia).




Figure 3. Antagonism test of rhizosphere fungi against *Alternaria alternata* on the 7th day. (note: (a): WO1; (b): WO2; (c): WO3; (d): WO4; (e): WO5; (f): WO6; (g): WO7; (h): WO8 ; (i): WO9; (j): WO10; (k): WO11; (l): WO12; (m): WO13; (n) control; A: Antagonistic fungi; P: Pathogenic fungi)




Figure 4. Mycoparasitism by WO9 (a1. attachment and coiling; a2. penetration; a3. lysis), and WO13 (b1. attachment; b2. penetration; b3. lysis); (Remarks: 1: antagonistic fungal hyphae; 2: pathogenic fungal hyphae)

Article in Progress