

Enhanced *Chlorella vulgaris* Buitenzorg growth by Photon Flux Density Alteration in Serial Bubble Column Photobioreactors

Anondho Wijanarko

Dianursanti Antonius Yudi Sendjaya

Misri Gozan

Roekmijati Widaningroem Soemantojo

Department of Chemical Engineering, Faculty of Engineering, The University of Indonesia
Kampus Universitas Indonesia, Depok 16424, INDONESIA

Arief Budi Witarto

Biotechnology Research Center, Indonesian Institute of Science, Jalan Raya Bogor KM 46,
Bogor 16911, INDONESIA

Kazuhiro Asami Kazuhisa Ohtaguchi

Department of Chemical Engineering, Tokyo Institute of Technology, 2-12-1 Ookayama,
Meguro ku, Tokyo 152, JAPAN

Micro algae are photolithotrophs that perform oxygenic photosynthesis and capable of accumulating a large amount of CO₂, using an inducible CO₂ concentrating mechanism (CCM). These characteristics make the micro algae potentially useful for removal and utilization of CO₂ emitted from industrial plant. Generally, the usage of photosynthetic microorganism in CO₂ fixation and biomass production for the economically viable commodities have been increased and significantly improved as a solution for this problem. Using these facts and previous research results using *Anabaena cylindrica* IAM M1 and *Spirulina platensis* IAM M 135, enhancement of CO₂ fixation and biomass production by *Chlorella vulgaris* Buitenzorg with photon flux density alteration along with an increasing of culture biomass during the cellular growth period, was implemented in this research. The photon flux density used in this alteration was the maximum light for *Chlorella's* maximum growth rate ($I_{m_{max, opt}}$). The cultivation of *Chlorella vulgaris* Buitenzorg in the Benneck basal medium operating conditions: T, 29°C; P, 1.0 atm; U_G, 2.4m/h; CO₂, 10%; using Philip Halogen Lamp 20W/12V/50Hz as the light source and three bubble column photobioreactors arranged in series order with each having a volume of 0.200dm³. Results had shown that the photon flux density alteration as a whole could increase around 60% the biomass production of *Chlorella vulgaris* and around 7% the CO₂ fixation ability, compared to constant photon flux density outcomes. This experiment also showed that the noncompetitive inhibition of [HCO₃⁻] as carbon source substrate is affected significantly during the cultivation in both of alteration and continuous photon flux density.

Keywords: Biomass, *Chlorella vulgaris* Buitenzorg, alteration, photobioreactor, seri

INTRODUCTION

Micro algae are photolithotrophs that perform oxygenic photosynthesis. Like higher plants, these organisms require radiant energy using water as an electron donor and represent significant contribution to eliminate atmospheric greenhouse gas carbon dioxide (CO₂) by ribulose biphosphate carboxylase/oxygenase (RubisCO). The reduced products of CO₂ serve as the carbon source for the entire biomaterial. In addition, micro algae are capable of accumulating a large amount of CO₂, using an inducible CO₂ concentrating mechanism (CCM) [Kaplan, Bdger, and Berry1980]. Previous study has shown that when micro algae *Synechococcus leopoliensis* IAM M6 was grown on the modified Detmer medium (MDM) at 313K, it accumulated external inorganic carbon ($c_i = \text{CO}_2 + \text{HCO}_3^-$) to the extent of 0.128 g.L⁻¹.h⁻¹ that was a five- to nine-fold higher than the CO₂ fixation rate of photosynthesis (Ohtaguchi, 2000). These characteristics make the micro algae potentially useful for study on the removal and utilization of CO₂ emitted from industrial plant to minimize the accumulation of carbon dioxide in nature. Generally, the usage of photosynthetic microorganisms *Anabaena cylindrica*, *Chlorella vulgaris*, *Synechococcus leopoliensis*, and *Spirulina platensis* in CO₂ fixation and biomass production for economically commodities such as has been increased and significantly improved; as a solution for this problem [Ohtaguchi and Wijanarko 2002; Wijanarko, Asami, and Ohtaguchi 2004a; Wijanarko and Ohtaguchi, 2004b].

Chlorella biomass is widely known as a high-potential substance with high economic values, such as chlorophyll, CGF, beta-carotene, protein and cell walls (Wirosapurto 2002). For its growth, *Chlorella* needs light energy and substrates. Light energy is an important factor for *Chlorella* growth that would be converted to ATP synthesizes for use in their photosynthesis, metabolism, growth, and cell division. Either,

the CO₂ as *Chlorella* substrate was fixated and used with the ATP synthesise in dark reaction to produced carbon compound (Ogbonna, 1995). Unfortunately, the results of CO₂ fixation and biomass production with constant photon flux density were relatively small, because the effect of self-shading phenomenon and the needs of light energy's increasing during microbial growth were considered as being slight [Falkowsky and Owens 1980; Oquist et al. 1992].

Using to these facts and the previous research results using *A. cylindrica* IAM M1 and *S. platensis* IAM M 135 (Wijanarko and Ohtaguchi 2003; Hirata, Taya, and Tone 1996, 1998), an enhancement effort of CO₂ fixation and biomass production by *C. vulgaris* Buitenzorg with photon flux density alteration along with an increasing of culture biomass during cellular growth period was implemented in this research. The photon flux density used in this alteration was the maximum light for *Chlorella*'s maximum growth rate $I_{m_{max,opr}}$.

This experimental report also contained the investigation results about the kinetic studies of microbial growth.

MATERIALS AND METHODS

C. vulgaris Buitenzorg was supplied from the Depok Research Center of Fresh Water Fishery, Indonesia. This strain was cultivated in a serial configuration of 0.200 dm³ bubble column photo bioreactor (0.08m inside (s), Pyrex glass made) using Benneck basal medium at T, 29°C; P, 1 atm; and, Q_G, 25.4 dm³/h. The inlet CO₂ composition enriched air $y_{CO_2,i}$ was set around 10% and illuminated by side illumination at 14.9W/m² (74.4 μmol/[m² s]) simultaneously until 103W/m² (542 μmol/[m²s]) refers to the increasing of biomass production at cultivation period with Philip Halogen lamp 20W/12V/50Hz. Cultivation at constant photon flux density (I_p) of 14.9W/m² (74.4μmol/[m² s]) was also performed as control experiment. Benneck was chosen as the medium since

it has a simple composition without carbon source and common nutrition for *C. vulgaris*'s development. Preculture conditioning was done in the beginning by feeding the air into the reactor and with the photon flux density at $2,95 \text{ W/m}^2$ ($14.7 \mu\text{mol}/[\text{m}^2 \text{ s}]$) in order to pass the lag phase of *C. vulgaris*'s growth. The photon flux density alteration performed a change of distance between one side of the photobioreactor and the parallel banks of halogen lamps as illumination source as shown in Figure 1.

Measuring OD_{600} of cell suspensions was performed for the growth analysis. The calibration curve is used to show that the value of OD_{600} is equivalent to the dry weight concentration. The CO_2 concentration of input and output gas ($y_{\text{CO}_2,i}$, $y_{\text{CO}_2,e}$) was performed by gas chromatography (GC TCD Shimadzu GC-8A). The photon flux densities of the incident and transmitted light were measured by Lux meter (*Luxtron LX-103*) and the pH of the cultures was measured using pH meter (*Hanna Model HI 8014*). Data




Figure 1. Experimental Apparatus

obtained from the experiment is processed to find several parameters as follows:

1. Biomass Amount (X)

The amount of cell biomass can be obtained using the calibration curve OD_{680} vs X .

2. The specific growth rate and incident specific growth rate (μ_{max} , μ) is calculated using the Monod's equation (Monod, 1942) as follow

$$\mu = \frac{1}{X} \cdot \frac{dX}{dt} \quad (1)$$

3. The carbon dioxide transfer rate (CTR) and specific carbon dioxide transfer rate (q_{CO_2}) is calculated using the following equation:

$$CRT = \Delta y_{CO_2} \cdot \alpha_{CO_2} \quad (2)$$

$$q_{CO_2} = \frac{CTR}{X} \quad (3)$$

$$\alpha_{CO_2} = Q_G \frac{M_{CO_2} \cdot P_T}{V \cdot R \cdot T} \quad (4)$$

Here, Q_G is gas volumetric rate, V is bioreactor volume, M_{CO_2} is molecular weight of CO_2 , P_T is ambient pressure, and T is ambient temperature.

4. The bicarbonate concentration ($[HCO_3^-]$) is calculated using Handerson-Hasselberg's equation as follows:

$$[HCO_3^-] = \frac{K_{CO_2,o} \cdot y_{CO_2} \cdot P_T}{H_{CO_2,o} \cdot 10^{-pH}} \cdot \left(\frac{\text{Exp} \left(A_K \cdot \left(1 - \frac{T_o}{T} \right) + B_K \cdot \ln \left(\frac{T}{T_o} \right) + C_K \cdot \left(\frac{T_o}{T} - 1 \right) \right)}{\text{Exp} \left(A_H \cdot \left(1 - \frac{T_o}{T} \right) + B_H \cdot \ln \left(\frac{T}{T_o} \right) + C_H \cdot \left(\frac{T_o}{T} - 1 \right) \right)} \right) \quad (5)$$

Here, $K_{CO_2,o}$ and $H_{CO_2,o}$ are equilibrium and Henry constant at 25°C and 1.0 atm., y_{CO_2} is ambient CO_2 concentration at ambient air, P_T is ambient pressure, pH is incident culture pH value. A_K , B_K , and C_K are thermodynamic equilibrium constant parameters while A_H , B_H , and C_H are thermodynamic Henry constant parameters.

5. The light energy utility for biomass production (E_x) was calculated by the following equations:

$$E_x = \frac{\int_0^t I_T \times dt}{X \times S} \quad (6)$$

Three equations of kinetic models were used to investigate the microbial growth model in this research as a substrate inhibition model (Bailey and Ollis 1986; Schugerl and Belgardt 2000), such as:

a. Monod

$$\mu = \mu_{max} \cdot \frac{[HCO_3^-]}{K_S + [HCO_3^-]} \quad (7)$$

b. Ierusalemky

$$\mu = \mu_{max} \cdot \frac{[HCO_3^-]}{K_S \cdot [HCO_3^-]} \cdot \frac{1}{\left(1 + \frac{[HCO_3^-]}{K_i} \right)} \quad (8)$$

c. Haldane

$$\mu = \mu_{max} \cdot \frac{[HCO_3^-]}{K_S + [HCO_3^-]} \cdot \frac{1}{\left(1 + \frac{[HCO_3^-]}{K_i} \right)} \quad (9)$$




Figure 2. Relationship between X and $I_{\mu_{max,opt}}$ of *C. vulgaris* Buitenzorg. Compared data is obtained from cultivation of *C. vulgaris* in the continuous photon flux density of 74.4 μmol/[m²s] (14,9 W/m²).

Here K_i and K_s are the inhibition and activation substrate constants.

RESULTS AND DISCUSSION

General Discussion

The biomass concentration used for initial running in each photobioreactor arranged in serial configuration is 0.81 g/dm³ of dry weight. This amount is an ideal low concentration that makes it have a possible to obtain a high specific growth rate. Cultivation started when the CO₂ gas flow was opened at a concentration of 10%. In the beginning of cultivation, the photon flux density given to the reactor was 14.9 W/m² (74.4 μmol/[m² s]). This is the optimum photon flux density ($I_{\mu_{max,opt}}$) for the cultivation of *C. vulgaris* which has biomass concentration was 0.81 g/dm³. Then, the photon flux density is increased during the cultivation period based on the amount of biomass inside each

photobioreactor. The photon flux density given to each reactor was based on the curve showing the relationship between the dry weight of cell (X) and the optimum photon flux density ($I_{\mu_{max,opt}}$) shown in Figure 2.

Experimental Data

From the cultivation of *Chlorella vulgaris* Buitenzorg in a serial bubble column photobioreactors with photon flux density alteration, the data is being shown out in Figure 3.

Meanwhile the control experiment results are shown in Figure 4.

Effect of Photon Flux Density Alteration to the Biomass Production of *C. vulgaris* Buitenzorg

Effect onto cellular dry weight (X). With the similar initial cellular dry weight of 0.81




Figure 3. Experimental results on photon flux density alteration during cell cultivation in serial bubble column photobioreactors

g/dm³, it was found that the growth curve's slope of *C. vulgaris* Buitenzorg is sharper in the photon flux density alteration compared with the constant photon flux density. This phenomenon happened in all of the photobioreactors in the serial configuration. The final biomass production result was tended by cellular dry weight after the cultivation entered the stationer phase as shown in Table 1.

Table 1. Final Cellular Dry Weight

Photon Flux Density	Reactor No.	X (g/dm ³)
ALTERATION	1	21.5
	2	19.5
	3	20.0
CONSTANT	1	8.2
	2	7.8
	3	8.2




Figure 4. Control Experiment Results by Constant Photon Flux Density during Cell Cultivation in Serial Bubble Column Photobioreactors

From Table 1, it is shown that the final biomass production results on the photon flux density alteration were more than the results on the constant photon flux density. These results tend that the photon flux density alteration was able to raise the cultivation of *C. vulgaris* Buitenzorg in biomass production by up to 60%. Hence, the photon flux density alteration based on cellular photosynthetic activation needs can reduce the self-shielding effect in dense cellular dry weight condition.

Effect onto the specific growth rate (μ). The values of specific growth rate (μ) during the

cultivation with photon flux density alteration and constant photon flux density are shown in the Figure 5.

Like the results of previous experiments (Wijanarko and Ohtaguchi 2003, Wijanarko et al. 2006), Figure 5 shows that photon flux density alteration increases the specific growth rate of the micro algae in the serial bubble column photobioreactors. This result was directly impacted to the amount of produced biomass.

Effect onto $[HCO_3^-]$. The culture's $[HCO_3^-]$ showed the amount of formed bicarbonate ions during cellular CO_2 fixation that was consumed




Figure 5. Specific Growth Rate (μ) Results in Serial Bubble Column Photobioreactors on (a) photon flux density alteration; and (b) constant photon flux density


by *Chlorella vulgaris* Buitenzorg during cellular growth. $[\text{HCO}_3^-]$ is calculated from the changing of the culture's pH as a result of the CO_2 fixation and of the cell growth activity of *C. vulgaris* Buitenzorg.

The carbon dioxide gas's entering the culture and hydrolyzing with water to form a bicarbonate ion was based on Eq. (10).



During the cellular growth period, *C. vulgaris* Buitenzorg cells consumed bicarbonate ion and, together with another culture's essential substrates, was metabolized by enzymatic reaction to form cellular building block organic compounds, such as: amino acid, fatty acid, glucose, sucrose, amyllum, cellulose, lignin, protein, fat, and oligo-protein.

Eq. 11 showed an overall enzymatic cellular HCO_3^- consumed reaction of *C. vulgaris* Buitenzorg to form glucose.



Using the approximation of Henry's Law and Hendersen-Hasselbach equation, the HCO_3^- concentration could be found from experimental pH data. Figure 6 shows the experimental HCO_3^- concentration of *C. vulgaris* Buitenzorg.

Because of intracellular growth activity, the culture pH increased and was directly impacted to the increased culture's $[\text{HCO}_3^-]$. Figure 6 shows that, at the end of running period, in each bubble column photobioreactor the photon flux density alteration mode provided 4.5mM of $[\text{HCO}_3^-]$, while the constant photon flux density mode tended 3.5mM. Thus, it was shown that the photon flux density alteration mode could enhance the culture's bicarbonate concentration up to 30%.

Effect of Photon Flux Density Alteration to *C. vulgaris* Buitenzorg's CO_2 Fixation

CTR showed the amount of transferred CO_2 gas to the culture medium during the cellular growth period. Figure 7 shows both of the CTR experimental results. Figure 7 shows




Figure 6. Cellular $[HCO_3^-]$ in Serial Bubble Column Photobioreactors on (a) photon flux density alteration and (b) constant photon flux density

that the average value of CTR of the photon flux density alteration (14.4 g/[dm³.h], equal to 1.38 tons CO₂/[ha.d], for a pond that has a maximum light path length of 40cm) is higher than the constant photon flux density (13.8 g/[dm³.h], equal to 1.30 tons CO₂/[ha.d]). Like the result for cellular day weight, the photon flux density alteration mode enhanced the value of CTR only up to around 5%. Compared with up to grass plants, such as *Mimosa* (0.133 tons CO₂/[ha.d]),

C. muconoides (0.121 tons CO₂/[ha.d]), and *Centronema* (0.296 tons CO₂/[ha.d]), the CO₂ fixated capability of *C. vulgaris* Buitenzorg in both photon flux density alteration and constant photon flux density is extremely higher than any of those plants capability (Widjaja 2002).

The q_{CO_2} showed the incident specific CO₂ fixation rate that was caused by biological activities during the cellular growth period. The value of q_{CO_2} can be defined as CTR divided by




Figure 7. CTR values in serial bubble column photobioreactors on (a) photon flux density alteration; (b) constant photon flux density.




Figure 8. q_{CO_2} Values in Serial Bubble Column Photobioreactors on (a) photon flux density alteration and (b) constant photon flux density.

the cellular dry weight. Figure 8 shows the two q_{CO_2} experimental results.

Figure 8 shows that the average value of q_{CO_2} of the photon flux density alteration (5.01 h^{-1}) is slightly higher than the constant photon flux density (4.92 h^{-1}) and that the photon flux density alteration mode enhances the value of q_{CO_2} only up to around 1%.

Energy for Biomass Production of *C. vulgaris* Buitenzorg.

Table 2 shows the calculated amount of utilized energy, which was used in the biomass production of *C. vulgaris* Buitenzorg.

Table 2. Energy for Biomass Production of *C. vulgaris* Buitenzorg

Photon Flux Density	Reactor No.	E_x (J/g)
ALTERATION	1	0.57
	2	0.81
	3	0.65
CONSTANT	1	1.66
	2	1.87
	3	1.67

Like in the results of previous experiment (Wijanarko et al. 2006), the value of average utilized energy on the photon flux density alteration mode is lesser than on constant photon flux density. The E_x on the photon flux density alteration is around 40% of the value on the constant photon flux density.

Kinetics Model of Cellular Substrate Consumption

Using the culture's $[HCO_3^-]$ as essential substrate for cellular growth, the kinetics curve fitting of Monod, Ierusalemky, and Haldane model equation approximation was carried out. This curve fitting is the method to figure out the matched model approximation which was able to describe the substrate consumption during the cultivation period. Figure 9 shows a matched model approximation curve fitting for the photon flux density alteration mode and Figure 10 for the constant photon flux density mode.

Although both the photon flux density alteration and the constant photon flux density modes showed that the activation and inhibition growth model of the Ierusalemky and Haldane equations have a relatively similar tendency. With reference to the previous experimental model results for filamentous cyanobacterium




Figure 9. Curve Fitting Result of Model Approximation of (a) Monod (b) Ierusalimsky, and (c) Haldane for Photon Flux Density Alteration




Figure 10. Curve Fitting Result of Model Approximation of (a) Monod; (b) Ierusalimsky, and (c) Haldane for Constant Photon Flux Density

Anabaena cylindrica Lemmerman on photon flux density alteration and using both single and serial reactors at low photon flux density of 1.0klx (Wijanarko and Ohtaguchi, 2003, 2004) and for unicellular green algae *C. vulgaris* Buitenzorg on photon flux density alteration in a single reactor at ambient temperature (Wijanarko et al. 2006), the Haldane equation is properly matched to the cellular growth kinetic of *C. vulgaris* Buitenzorg on both photon flux density alteration and constant photon flux density in a serial reactor. This means that in both modes carbon source significantly affects in the competitively inhibition mechanism of Haldane.

CONCLUSION

Alteration of light illumination in *Chlorella vulgaris* Buitenzorg cultivation can increase the biomass production (X) by 60% higher than biomass production in continuous illumination when photon flux density is $I_{\mu_{max,opt}}$ at the basic inoculum.

The capacity of *C. vulgaris* Buitenzorg for CO₂ fixation slightly increases to around 5% with that in photon flux density alteration compare to the constant photon flux density.

The utilized energy for biomass production of *C. vulgaris* Buitenzorg on the photon flux density alteration is less than constant photon flux density, by around 40% of the value of constant photon flux density.

The kinetics equation model of Haldane is most satisfactory for biomass production on both photon flux density alteration and constant photon flux density. It means that in both modes, carbon source is significantly affecting in the competitively inhibition mechanism.

ACKNOWLEDGEMENT

The authors thank the reviewers and editors for their useful comments on this paper. This work was supported in part by 13th Competitive Grant for Scientific Research of the Directorate General of

Higher Education, Ministry of National Education, Republic of Indonesia.

REFERENCES

- Bailey, J.E. and Ollis, D.F. (1986), *Biochemical Engineering Fundamental*, McGraw Hill Book Co., New York Ch.7, 382-392.
- Falkowsky, P.G. and Owens, T.G. (1980), "Light-Shade Adaptation," *Plant Physiology*, 66, 592-595.
- Hirata, S., Taya, M. and Tone, S. (1996). "Characterization of *Chlorella* cell cultures in batch and continuous operations under a photoautotrophic condition," *Journal of Chemical Engineering of Japan*, 29, 6, 953-959.
- Hirata, S., Taya, M. and Tone, S. (1998). "Continuous cultures of *Spirulina platensis* under photoautotrophic conditions with change in light intensity," *Journal of Chemical Engineering of Japan*, 31, 4, 636-639.
- Kaplan A., Badger, M.R. and Berry, J.A. (1980), "Photosynthesis and the intracellular inorganic carbon pool in the blue green alga *Anabaena variabilis*: Response to external CO₂ concentration," *Planta*, 149, 219-226.
- Monod, J. (1942), *Recherche sur la croissance des cultures bacteriennes*. Herman and Cie, Paris, 211-232.
- Ogbonna, J. C., Yada, H. and Tanaka, H. (1995). "Kinetics study on light-limited batch cultivation of photosynthetic cells", *Journal Fermentation and Bioengineering*, 80, 7, 259-264.
- Ohtaguchi, K. (2000). "Soft energy path synthesis from carbon dioxide to biofuel ethanol through cyanobacterial biotechnology," *Technology*, 7S, 175-188.
- Ohtaguchi, K. and Wijanarko, A. (2002). "Elevation of the efficiency of cyanobacterial carbon dioxide removal by monoethanolamine solution," *Technology*, 8, 267-286.

- Oquist, G., Anderson, J.M., McCaffery, S. and Chow, W. S. (1992). "Mechanistic difference in photoinhibition of sun and shade plants," *Planta*, 188, 422-431.
- Schugerl, K. and Bellgardt, K.H. eds. (2000). *Bioreaction Engineering Modeling and Control*, Springer, Berlin.
- Widjaja, H. (2002). *Penyimpanan karbon dalam tanah, alternatif carbon sink dari pertanian konservasi, Diktat Pengantar Falsafah Sains*, Bogor Institute of Agriculture Press, 1-14.
- Wijanarko, A. and Ohtaguchi, K. (2003). "Alteration of light illumination during microbial growth: An enhancement effort of biomass production and carbon dioxide fixation of psychrophilic cyanobacterium *Anabaena cylindrica* IAM M1," *Comparative Biochemistry and Physiology*, 134, 124.
- Wijanarko, A. and Ohtaguchi, K. (2004). "Carbon dioxide removal and biomass production by *Anabaena cylindrica* IAM M1 using reactor in series", *Study on Surface Science and Catalysis*, 153, 461-468.
- Wijanarko, A., Asami, K. and Ohtaguchi, K. (2004). "The kinetics of growth and the CO₂ concentrating mechanism of the filamentous cyanobacterium *Anabaena cylindrica* in a bubble column," *Journal of Chemical Engineering of Japan*, 37, 8, 1019-1025.
- Wijanarko, A., Dianursanti, H., R.W. Soemantojo, and Ohtaguchi, K. (2006). "Effect of light illumination alteration on *Chlorella vulgaris* Buitenzorg's CO₂ fixation in bubble column photobioreactor," **International Journal on Algae**, 8, 1, 53-60.
- Wirosaputro, S. (2002), *Chlorella Untuk Kesehatan Global*, Gadjah Mada University Press, Yogyakarta, Indonesia.
-